

Symmetries of the Robinson–Trautman equation

W. Natorf and J. Tafel,
Institute of Theoretical Physics,
Warsaw University, ul. Hoża 69 00–681,
Warsaw, Poland

Abstract

We study point symmetries of the Robinson–Trautman equation. The cases of one- and two-dimensional algebras of infinitesimal symmetries are discussed in detail. The corresponding symmetry reductions of the equation are given. Higher dimensional symmetries are shortly discussed. It turns out that all known exact solutions of the Robinson–Trautman equation are symmetric.

1 Introduction

In 1960 Robinson and Trautman introduced¹ a class of space-times admitting a diverging shear- and twist-free congruence of null geodesics. Such space-times, if asymptotically Minkowskian, are believed to describe gravitational radiation outgoing from spatially bounded sources. Recent numerical results suggest that the Robinson–Trautman metrics can be used to estimate the mass loss during the final phase of the collision of two black holes.² There are also suggestions that the subclass of the so called C-metrics³ (and their twisting generalizations⁴) can describe spacetimes containing accelerating black holes.^{5–8}

In terms of standard coordinates $u, r, \xi, \bar{\xi}$, where u and r are real and ξ is complex, the Robinson–Trautman metrics are given by⁹

$$g = 2du(Hdu + dr) - 2r^2 P^{-2} d\xi d\bar{\xi} . \quad (1)$$

The function P is independent of r . Vacuum Einstein equations imply

$$H = -r\partial_u \ln P - m(u)/r + P^2 \partial_\xi \partial_{\bar{\xi}} \ln P \quad (2)$$

and a fourth order equation for P , referred to as the Robinson–Trautman equation:

$$P^2 \partial_\xi \partial_{\bar{\xi}} (P^2 \partial_\xi \partial_{\bar{\xi}} \ln P) + 3m\partial_u (\ln P) - \partial_u m = 0 . \quad (3)$$

Using coordinate freedom m can be transformed to the value ± 1 or 0 . The Gaussian curvature K of surfaces of constant u and r is given by

$$K = 2P^2 \partial_\xi \partial_{\bar{\xi}} \ln P . \quad (4)$$

Existence of asymptotically flat Robinson–Trautman metrics has been recently examined.^{10–12} Also, their large u asymptotic behaviour is known.^{13,14} These results, however, do not give any hint of how to look for explicit solutions. Only a few of them are known since 1960, none of them being asymptotically flat except for the cases of the Minkowski and the Schwarzschild metrics.

Assuming asymptotic flatness of metrics (1), the Bondi energy,^{15,16} the Bondi mass aspect and the news function were found in terms of P .^{11,17–19} Asymptotic flatness of (1) with positive m follows from the assumption that in the gauge $m = 1$ the function $\hat{P} = P/(1 + \frac{1}{2}\xi\bar{\xi})$ is positive and regular on $R \times S_2$, ξ being interpreted as a complex stereographic coordinate on S_2 .

The subclass of metrics with $m = 0$ is characterized by the fact that K is a solution of the Laplace equation, so K must be either constant or singular. No nontrivial asymptotically flat metric exists in this subclass.

In the present paper we consider symmetries of the Robinson–Trautman equation and classify the conjugacy classes of one- and two-dimensional symmetry algebras. We find symmetry reductions of equation (3) for solutions preserved by these algebras. Some of these solutions might correspond to asymptotically flat metrics with a simple dependence on the time u . This research may also be of help for studying the numerical solutions of the Robinson–Trautman equation, providing natural ansatzes with smaller number of variables than in generic situations.

2 Symmetry transformations for $m \neq 0$

Suppose $m \neq 0$ and consider equation (3) in the gauge $m = 1$,

$$P^2 \partial_\xi \partial_{\bar{\xi}} (P^2 \partial_\xi \partial_{\bar{\xi}} \ln P) + 3 \partial_u \ln P = 0 . \quad (5)$$

It is easy to prove that (5) is invariant with respect to the following point transformations:

$$\begin{aligned} u &\mapsto u' = a^4 u + b, \\ \xi &\mapsto \xi' = f(\xi), \\ P &\mapsto P' = a^{-1} |f_{,\xi}| P , \end{aligned} \quad (6)$$

where $a \neq 0$ and b are real constants and f is a holomorphic function of ξ and $f_{,\xi} \neq 0$. If (6) is supplemented by

$$r' = a^2 r , \quad (7)$$

the corresponding metric transforms as $g \mapsto a^6 g$. Thus, (6) together with (7) induces a homothety of the metric. It is an isometry when $a = 1$.

Infinitesimal transformations corresponding to (6) are given by the vector field

$$k = (4Au + B) \partial_u + F(\xi) \partial_\xi + \bar{F}(\bar{\xi}) \partial_{\bar{\xi}} + (\text{Re} F_{,\xi} - A) P \partial_P , \quad (8)$$

where A and B are real constants and F is a holomorphic function of ξ . These fields form the symmetry algebra $\mathfrak{g}$.

Solutions of (3) invariant with respect to (8) have to satisfy the following linear equation

$$(4Au + B) P_{,u} + F P_{,\xi} + \bar{F} P_{,\bar{\xi}} + (A - \text{Re} F_{,\xi}) P = 0 . \quad (9)$$

We will perform the symmetry reduction of (5) assuming that its solution is preserved by vector fields (8) forming a one- or two-dimensional subalgebra of $\mathfrak{g}$ (higher dimensional subalgebras will also be shortly discussed.) First we will classify these subalgebras under the action of pseudogroup of transformations (6). This way we will obtain the conjugacy classes (CC) of one- and two-dimensional subalgebras of $\mathfrak{g}$.

3 Solutions with one or two symmetries

Consider a one-dimensional algebra generated by a vector field (8). Applying an appropriate transformation (6) we can simplify the coefficients A and B and the function F , obtaining one representative of each conjugacy class. For instance, if $A = 0$, $B \neq 0$ and $F \neq 0$ we can scale u and ξ so that $B = 1$ and $F = \xi$. This leads to

$$k = \partial_u + \xi \partial_\xi + \bar{\xi} \partial_{\bar{\xi}} + P \partial_P . \quad (10)$$

In the same way we can distinguish five conjugacy classes which are listed in Table 1, together with corresponding vector k , form of the invariant solution P and the reduced Robinson–Trautman equation. Whenever it is needed, we explicitly write the definition of the new variable $z(u, \xi)$ appearing in the invariant solution. Throughout the text we use x and y to denote, respectively, the real and imaginary part of ξ .

Consider now the case of two-dimensional subalgebra $\mathfrak{g}_2$ of $\mathfrak{g}$. We denote the basis vectors of $\mathfrak{g}_2$ by k_1 and k_2 . There are two nonisomorphic two-dimensional Lie algebras such that either

$$[k_1, k_2] = 0 \quad (11)$$

or

$$[k_1, k_2] = k_2 . \quad (12)$$

The Lie bracket of two fields given by (8) reads

$$\begin{aligned} [k_1, k_2] &= 4(B_1 A_2 - B_2 A_1) \partial_u + (F_1 F_{2,\xi} - F_2 F_{1,\xi}) \partial_\xi + \\ &\quad + (\bar{F}_1 \bar{F}_{2,\bar{\xi}} - \bar{F}_2 \bar{F}_{1,\bar{\xi}}) \partial_{\bar{\xi}} + P \text{Re}(F_1 F_{2,\xi\xi} - F_2 F_{1,\xi\xi}) \partial_P \end{aligned} \quad (13)$$

where the indices 1, 2 refer to vectors k_1, k_2 respectively.

In the Abelian case, equation (11) implies

$$A_1 B_2 = A_2 B_1 , \quad (14a)$$

$$F_1 F_{2,\xi} = F_2 F_{1,\xi} , \quad (14b)$$

$$\text{Re}(F_1 F_{2,\xi\xi} - F_2 F_{1,\xi\xi}) = 0 . \quad (14c)$$

It follows from (14b) that F_1 is proportional to F_2 and (14c) is satisfied.

Due to (14a) we can assume without loss of generality that $A_2 = B_2 = 0$. Then $F_2 \neq 0$ and using the gauge freedom in ξ we can set $F_2 = i$. Then $F_1 = C = \text{const} \in R$ follows from (14b) and the remaining freedom in choice of k_1 . Therefore, the symmetry generators are

$$k_1 = (4Au + B)\partial_u + C\partial_x - AP\partial_P, \quad k_2 = \partial_y . \quad (15)$$

Invariance of a solution P of (5) with respect to k_2 implies

$$P = p(u, x) . \quad (16)$$

Invariance with respect to k_1 gives

$$(4Au + B)P_{,u} + CP_{,x} = -AP . \quad (17)$$

Depending on values of A, B and C we can distinguish five conjugacy classes of two-dimensional Abelian subalgebras of $\mathfrak{g}$. They are listed in Table 2. (Here, as well as in Tables 3 and 4 the symbol p' denotes a derivative of p with respect to its argument.)

In the non-Abelian case, equation (12) gives

$$4(A_2 B_1 - A_1 B_2) = 4A_2 u + B_2 , \quad (18a)$$

$$F_1 F_{2,\xi} - F_2 F_{1,\xi} = F_2 , \quad (18b)$$

$$P \text{Re}(F_1 F_{2,\xi\xi} - F_2 F_{1,\xi\xi}) = P(\text{Re} F_{2,\xi} - A_2) . \quad (18c)$$

It follows from (18a) that

$$A_2 = 0, \quad B_2(1 + 4A_1) = 0 . \quad (19)$$

Differentiating (18b) and its complex conjugate we conclude that (18c) follows from (18b). Proceeding as in the Abelian case, we obtain results summarized in Table 3.

4 Solutions with more symmetries

All Lie algebras of dimension greater than three contain a three-dimensional subalgebra. Let $k_i, i = 1, 2, 3$, be its generators. In the case of Bianchi type *VIII* or *IX* it follows from (13) that $A_i = B_i = 0$. Then the fields k_1, k_2 and k_3 are tangent to a two-dimensional surface and invariant solutions are excluded. For all other Bianchi types the algebra contains a two-dimensional Abelian subalgebra. One can construct invariant solutions as special cases of those described in Table 2. After lengthy calculations one obtains only the trivial solution $P = \text{const}$ or $P = \text{const} \cdot x^{3/2}$ given by Robinson and Trautman.²⁰ These are also solutions of the Robinson–Trautman equation for $m = 0$. Thus, assumption of three or more symmetries does not lead to any new interesting solutions.

5 Symmetry transformations for $m = 0$

Suppose now that $m = 0$. In this case equation (3) can be integrated to a second order equation,

$$P^2 \partial_\xi \partial_{\bar{\xi}} \ln P = \operatorname{Re} \phi(u, \xi) , \quad (20)$$

where ϕ is holomorphic with respect to ξ . If $\phi, \xi \neq 0$, one can transform (20) to the equation⁹

$$P^2 \partial_\xi \partial_{\bar{\xi}} \ln P = -\operatorname{Re} \xi . \quad (21)$$

Substituting

$$P = x^{3/2} p(\xi, \bar{\xi}) \quad (22)$$

into (21) we get

$$p^2 (\Delta \ln p - 3/8) = -1 . \quad (23)$$

Here $\Delta = x^2 \partial_\xi \partial_{\bar{\xi}}$ is the Laplace operator on a pseudosphere with the metric

$$g = \frac{d\xi d\bar{\xi}}{x^2} \quad (24)$$

which can be put into the standard form

$$g = \frac{4d\zeta d\bar{\zeta}}{(1 - \zeta\bar{\zeta})^2} \quad (25)$$

by means of the transformation $\zeta = (\xi + 1)/(\xi - 1)$. The operator Δ is preserved by transformations

$$\xi \mapsto \xi' = \frac{a\xi + ib}{ic\xi + d} , \quad a, b, c, d \in R \quad (26)$$

corresponding to an action of $\operatorname{SL}(2, R)$ on the pseudosphere.

It follows from (26) that infinitesimal transformations are generated by vector fields of the form

$$k = (iA\xi^2 + B\xi + iC)\partial_\xi + \text{c.c.} , \quad (27)$$

where A, B and C are real constants. Using symmetry transformations we can distinguish three conjugacy classes of one-dimensional subalgebras of the symmetry algebra for $m = 0$ (Table 4).

Given a solution of (21) we can apply the transformation $\xi' = f(u, \xi)$ to obtain a class of solutions of (20).

Note that the only solution of equation (23) invariant with respect to two independent fields of the form (27) is $p = \sqrt{8/3}$ which gives the well known solution $P = \text{const} \cdot x^{3/2}$ found by Robinson and Trautman.²⁰

Table 1. Invariant solutions with single symmetry.

CC	k	P	RT equation
1	∂_u	$p(\xi, \bar{\xi})$	$p^2 \partial_\xi \partial_{\bar{\xi}} \ln p = \operatorname{Re} \xi.$
2	$4u\partial_u - P\partial_P$	$u^{-1/4}p(\xi, \bar{\xi})$	$p^2 \partial_\xi \partial_{\bar{\xi}} (p^2 \partial_\xi \partial_{\bar{\xi}} \ln p) = \frac{3}{4}.$
3	$\partial_u - \xi \partial_\xi - \bar{\xi} \partial_{\bar{\xi}} - P\partial_P$	$e^u p(z, \bar{z})$	$p^2 \partial_z \partial_{\bar{z}} (p^2 \partial_z \partial_{\bar{z}} \ln p) - 3(z \partial_z + \bar{z} \partial_{\bar{z}}) \ln p + 3 = 0, \quad z = e^{-u} \xi.$
4	∂_y	$p(u, x)$	$p^2 \partial_x^2 (p^2 \partial_x^2 \ln p) + 3 \partial_u \ln p = 0.$
5	$4Au\partial_u + \xi \partial_\xi + \bar{\xi} \partial_{\bar{\xi}} + P(1-A)\partial_P$	$u^{\frac{1-A}{4A}} p(z, \bar{z})$	$4Ap^2 \partial_z \partial_{\bar{z}} (p^2 \partial_z \partial_{\bar{z}} \ln p) + 3(A-1) + 3(z \partial_z + \bar{z} \partial_{\bar{z}}) \ln p = 0,$ $z = u^{-(4A)^{-1}} \xi.$

Table 2. Invariant solutions with two Abelian symmetries.

CC	k_1	k_2	P	RT equation
A.1	∂_u	∂_y	$p(x)$	$p^2 (\ln p)'' = x.$
A.2	∂_x	∂_y	$p(u)$	$p' = 0.$
A.3	$\partial_u + \partial_x$	∂_y	$p(u-x)$	$p^2 (p^2 (\ln p)'')'' + 3 (\ln p)' = 0.$
A.4	$4u\partial_u - P\partial_P$	∂_y	$u^{-1/4} p(x)$	$p^2 (p^2 (\ln p)'')'' = \frac{3}{4}.$
A.5	$4Au\partial_u + \partial_x - AP\partial_P$	∂_y	$u^{-1/4} p((4A)^{-1} \ln u - x)$	$p^2 (p^2 (\ln p)'')'' + \frac{3}{4A} (\ln p)' = \frac{3}{4}.$

Table 3. Invariant solutions with two non-Abelian symmetries.

CC	k_1	k_2	P	RT equation
NA.1	$\epsilon \partial_u - x \partial_x - y \partial_y - P \partial_P,$ $\epsilon = 0$ or $\epsilon = 1$	∂_y	$x p(u + \epsilon \ln x)$	$\epsilon[p^3(p'''' - 2p''' - p'' + 2p') -$ $-p^2(p''^2 + 2p'p'' + p'^2)] + 3(\ln p)' = 0.$
NA.2	$4Au \partial_u - x \partial_x - y \partial_y - (1 + A)P \partial_P$	∂_y	$u^{-\frac{1+A}{4A}} p(\underbrace{u^{\frac{1}{4A}} x}_X)$	$p^2(p^2(\ln p)'')'' + 3X(\ln p)' = 3(1 + A).$
NA.3	$-u \partial_u + \partial_x + \frac{1}{4}P \partial_P$	∂_u	$p(x)$	$p^2(\ln p)'' = x.$
NA.4	$-u \partial_u + \partial_x + \frac{1}{4}P \partial_P$	$\partial_u + e^\xi \partial_\xi + e^{\bar{\xi}} \partial_{\bar{\xi}} +$ $+P \operatorname{Re} e^\xi \partial_P$	$u^{-\frac{1}{4}} 1 + u^{-1} e^{-\xi} ^{\frac{3}{4}} p(z),$ $z = \frac{1+u^{-1} e^{-\xi}}{1+u^{-1} e^{-\bar{\xi}}}.$	$p^2(p^2(\ln p)'')'' + \frac{1}{4}p^4(\ln p)'' -$ $-36 \cos z + 48(\ln p)' \sin z = 0.$

Table 4. Invariant solutions for $m = 0$.

CC	k	P	RT equation
0.1	$i(\xi^2 - 1)\partial_\xi - i(\bar{\xi}^2 - 1)\partial_{\bar{\xi}}$	$p(r), r = \left \frac{\xi+1}{\xi-1} \right $	$\frac{1}{4}(1 - r^2)^2 \frac{1}{r}(r(\ln p)')' - \frac{3}{8} + p^{-2} = 0.$
0.2	$\xi \partial_\xi + \bar{\xi} \partial_{\bar{\xi}}$	$p(\phi), \xi = r e^{i\phi}$	$\frac{1}{4} \cos^2 \phi (\ln p)'' - \frac{3}{8} + p^{-2} = 0.$
0.3	$i(\partial_\xi - \partial_{\bar{\xi}})$	$p(x)$	$\frac{1}{4}x^2(\ln p)'' - \frac{3}{8} + p^{-2} = 0.$

Discussion

We have examined point symmetries of the Robinson–Trautman equation (3). Forms of solutions in the case of one or two symmetries were given, as well as the corresponding reduced equations. Note that all known⁹ exact solutions of (3) have two or three symmetries and belong to one of the cases considered here.

Assumption of symmetry does not seem to exclude regular solutions corresponding to asymptotically flat metrics. The coordinates $\xi, \bar{\xi}$ used in Tables 1–3 can differ from the hypothetical Bondi–Sachs coordinates. For instance, in the case 4 of table 1, the function P depends on x which becomes $\ln|\xi'|$ under the transformation $\xi \mapsto \xi' = \exp(\xi)$. To achieve regularity of $\hat{P}$ one should demand that P is everywhere finite, positive, and

$$\lim_{|\xi'| \rightarrow \infty} |\xi'|^{-1} P = a(u), \quad \lim_{|\xi'| \rightarrow 0} |\xi'| P = b(u) \quad (28)$$

for some positive functions a and b . Rewriting (28) in terms of ξ we get

$$\lim_{x \rightarrow -\infty} e^x P = a(u), \quad \lim_{x \rightarrow \infty} e^{-x} P = b(u) \quad (29)$$

which means that P behaves like $e^{|x|} f(u)$ for large $|x|$.

In the case A.3 (see Table 2) the Robinson–Trautman equation can be solved analytically. This way one obtains the so-called C-metrics. A possible physical interpretation of these metrics as well as their twisting generalizations (the spinning C-metrics) was done recently^{5–8}

Acknowledgments

J. T. would like to thank M. Grundland who participated in the early stages of these investigations.

This work was partially supported by the Polish Committee for Scientific Research (grant 1 PO3B 075 29).

References

- ¹Robinson I. and Trautman A., *Spherical Gravitational Waves*, Phys. Rev. Lett. **4** 461 (1960)
- ²Moreschi O., Perez A., and Lehner L., *Energy and angular momentum radiated for non head-on binary black hole collisions*, Phys. Rev. D **66** 104017 (2002)
- ³Kinnersley W., *Field of an Arbitrarily Accelerating Point Mass*, Phys. Rev. **186**, 1335 (1969)
- ⁴Demiański M., Plebański J. F., *Rotating, charged and uniformly accelerating mass in general relativity* Ann. Phys. (USA) **98** 98 (1976)
- ⁵Pravda V. and Pravdová A., *Boost-rotation symmetric vacuum spacetimes with spinning sources* J. Math. Phys. **43** 1536 (2002)
- ⁶Pravda V. and Pravdová A., *Boost-rotation symmetric spacetimes—review* Czech. J. Phys. **50** 333 (2000)
- ⁷Pravda V. and Pravdová A., *On the spinning C-metric* Gravitation: Following the Prague Inspiration (Selected essays in honour of J. Bičák), Eds. O. Semerak, J. Podolský and M. Zofka, World Scientific, Singapore (2002)
- ⁸Bičák J., Pravda V., *Spinning C-metric: a spacetime with accelerating, rotating black holes*, Phys. Rev. D **40** 1827 (1989)
- ⁹Stephani H., Kramer D., MacCallum M., Hoenselaers C., and Herlt E., *Exact Solutions to Einstein's Field Equations, Second Edition*, 2003, Cambridge University Press.
- ¹⁰Rendall A., *Existence and asymptotic properties of global solutions of Robinson–Trautman equation*, Class. Quantum Grav. **5** 1339 (1988)
- ¹¹Singleton D., *On global existence and convergence of vacuum Robinson–Trautman solutions*, Class. Quantum Grav. **7** 1333, (1990)

- ¹²Chruściel P. T., *On the global structure of Robinson–Trautman space-times*, Proc. R. Soc. Lond. A **436** 299 (1992)
- ¹³Foster J., and Newman E. T., *Note on the Robinson–Trautman Solutions*, J. Math. Phys. **8** 189 (1966)
- ¹⁴Frittelli S., and Moreschi O. M., *Study of the Robinson–Trautman metrics in the asymptotic future*, Gen. Rel. Grav. **24** 575 (1992)
- ¹⁵Bondi H., *Gravitational Waves in General Relativity*, Nature (London) **186** 535 (1960)
- ¹⁶Bondi H., van der Burg M. G. J., and Metzner A. W. K., *Gravitational Waves in General Relativity. VII. Waves from Axi-Symmetric Isolated Systems*, Proc. Roy. Soc. London **A269** 21 (1962)
- ¹⁷Tod P., *Some Examples of Penrose’s Quasi Local Mass Construction*, Proc. Roy. Soc. Lond. A **388** 457, (1983)
- ¹⁸Tafel J., *Bondi mass in terms of the Penrose conformal factor*, Class. Quantum Grav. **17** 4397 (2000)
- ¹⁹Cornish F. H. J., and Micklewright B., *The news function for Robinson–Trautman radiating metrics* Class. Quantum Grav. **16** 611 (1999)
- ²⁰Robinson I., and Trautman A., *Some spherical gravitational waves in general relativity*, Proc. Roy. Soc. London A **265** 463 (1962)